

DIGESTIVE DISEASES CENTER OF SOUTH TEXAS, P.L.L.C.

K.P. GANESHAPPA, M.D., F.A.C.G., F.A.C.P.
RAVI BOTLA, M.D., F.A.C.G.

HECTOR D. ALLENDE, M.D., F.A.C.P.
ERIC LAWITZ, M.D., C.P.I.

RAVI GANESHAPPA, M.D.

DIPLOMATES, AMERICAN BOARD OF INTERNAL MEDICINE AND GASTROENTEROLOGY

Cocktail Morning Procedure

Laxative Prep Instructions

ON THE DAY BEFORE THE EXAM you will need to be on a clear liquid diet the entire day. Examples of Clear Liquids include: Fat-Free Broth, Bouillon, Liquid portions of soup, Coffee, Tea, Pulp-Free juices, Sodas, Popsicles, etc. One or two ounces of milk to flavor your coffee are OK to use. Please AVOID anything with BRIGHT artificial colorings such as RED, BLUE, or GREEN dye (this will turn your colon that color!) AVOID JELL-O as this will harden your stools. It is important to drink adequate fluids throughout the day. **DO NOT EAT OR DRINK ANYTHING AFTER MIDNIGHT.**

At 5 PM (you can start earlier if you want) you will begin by mixing 2 oz. of Lactulose, 1 Capful of Miralax (17 g) in 8oz of Orange Juice, Apple Juice, Ginger Ale, Ice Tea, Lemonade, Sprite, or Water. If you are a diabetic use Water, Diet Sprite, Diet Ginger Ale, or Non-Sweetened ice Tea. Take this mixture every 30 – 45 minutes for a total of 6 doses. If you get too full, you may spread the doses further apart (ex, you can take the doses every hour instead of every 45 minutes). You can drink as much water as you want during this time. If you have heart failure or are on dialysis, try to limit the amount of water you drink.

Medications

1. Stop taking Coumadin (a blood thinning medication) and Iron 7 days before the procedure, if possible. Please consult with your primary care physician to make sure it is OK to stop your Coumadin. We will check your blood tests the day before the procedure to make sure that your blood is not too thin.
2. Please let us know if you are taking any of the following: Aspirin, Plavix, Bufferin, Mortin, Ibuprofen, Advil, Naprosyn, Aleve, Nuprin, and Alka-Seltzer.

Diabetic Patients

1. Do not take oral diabetic medication (Pills) the day of the exam.
2. If Insulin is used at bedtime, take half of your usual dose.
3. If Insulin is used twice a day, take half of your morning dose on the day of the exam.

Patient Name: _____ - **Date of Exam:** ____/____/____

Time of Exam: _____ A.M / P.M - **Arrival Time:** _____ A.M / P.M - **Facility:** _____

Please make arrangements for someone to drive you home after the exam. Due to the medication administered to sedate you, you will be unable to drive or take on any responsible tasks for the remainder of the day. The exam WILL NOT BE DONE IF YOU COME BY YOURSELF, VIA BUS, OR TAXI CAB.

You should be ready for discharge approximately 30 minutes after the exam.

There is a possibility that your insurance company may not cover these procedures, we suggest that you contact your carrier prior to the exam to verify coverage. All unpaid charges will be the responsibility of the patient.


497 10TH ST., SUITE 102
FLORESVILLE, TEXAS 78114
TELEPHONE: (210) 253-3422

621 CAMDEN STREET
SAN ANTONIO, TEXAS 78215
TELEPHONE: (210) 253-3422

1901 HWY 97E, SUITE 200
JOURDANTON, TX 78026
TELEPHONE: 830-769-5259

TOLL FREE #: 1-800-278-2571

WWW.DIGDISCTR.COM



DIGESTIVE DISEASES CENTER OF SOUTH TEXAS, P.L.L.C.

K.P. GANESHAPPA, M.D., F.A.C.G., F.A.C.P.
RAVI BOTLA, M.D., F.A.C.G.

HECTOR D. ALLENDE, M.D., F.A.C.P.
ERIC LAWITZ, M.D., C.P.I.

RAVI GANESHAPPA, M.D.

DIPLOMATES, AMERICAN BOARD OF INTERNAL MEDICINE AND GASTROENTEROLOGY

497 10TH ST., SUITE 102
FLORESVILLE, TEXAS 78114
TELEPHONE: (210) 253-3422

621 CAMDEN STREET
SAN ANTONIO, TEXAS 78215
TELEPHONE: (210) 253-3422

1901 HWY 97E, SUITE 200
JOURDANTON, TX 78026
TELEPHONE: 830-769-5259

TOLL FREE #: 1-800-278-2571

WWW.DIGDISCTR.COM